arXiv:1004.1245v1 [math.GR] 8 Apr 2010

MSC2010 20D20, 20E34

A conjugacy criterion for Hall subgroups in finite groups*

E.P. Vdovin, D.O. Revin

April 9, 2010

Abstract

A finite group G is said to satisfy C_π for a set of primes π, if G possesses exactly one class of conjugate π-Hall subgroups. In the paper we obtain a criterion for a finite group G to satisfy C_π in terms of a normal series of the group.

Introduction

Let π be a set of primes. We denote by π' the set of all primes not in π, by $\pi(n)$ the set of prime divisors of a positive integer n, while for a finite group G by $\pi(G)$ we denote $\pi(|G|)$. A positive integer n with $\pi(n) \subseteq \pi$ is called a *π-number*, while a group G with $\pi(G) \subseteq \pi$ is called a *π-group*. A subgroup H of G is called a *π-Hall subgroup*, if $\pi(H) \subseteq \pi$ and $\pi(|G : H|) \subseteq \pi'$. According to [1] we say that G *satisfies* E_π (or briefly $G \in E_\pi$), if G possesses a π-Hall subgroup. Moreover, if every two π-Hall subgroups are conjugate, then we say that G *satisfies* C_π ($G \in C_\pi$). Further if each π-subgroup of G lies in a π-Hall subgroup then we say that G *satisfies* D_π ($G \in D_\pi$). A group satisfying E_π (respectively, C_π, D_π) we also call an E_π- (respectively, C_π-, D_π-) *group*.

Let A, B, and H be subgroups of G such that $B \trianglelefteq A$. We denote $N_H(A) \cap N_H(B)$ by $N_H(A/B)$. Then every element $x \in N_H(A/B)$ induces an automorphism of A/B acting by $Ba \mapsto Bx^{-1}ax$. Thus the homomorphism $N_H(A/B) \to \mathrm{Aut}(A/B)$ is defined. The image of the homomorphism is denoted by $\mathrm{Aut}_H(A/B)$ and is called the *group of H-induced automorphisms on A/B*, while the kernel is denoted by $C_H(A/B)$. If $B = 1$, then $\mathrm{Aut}_H(A/B)$ is denoted by $\mathrm{Aut}_H(A)$.

Assume that π is fixed. It is proved that the class of D_π-groups is closed under homomorphic images, normal subgroups (mod CFSG[1]), [2, Theorem 7.7] or [3, Corollary 1.3]), and extensions (mod CFSG, [2, Theorem 7.7]). Thus, a finite group G satisfies D_π if and only if each composition factor S of G satisfies D_π. The class of E_π-groups is also known

---

*The authors were supported by Russian Foundation for Basic Research (Grants 08-01-00322 and 10-01-00391), the Russian Federal Agency for Education (Grant 2.1.1.419), and Federal Target Grant (Contract No. 02.740.11.0429). The first author gratefully acknowledges the support from Deligne 2004 Balzan prize in mathematics, and the Lavrent'ev Young Scientists Competition (No 43 on 04.02.2010).

[1] (mod CFSG) in the paper means that the result is proven modulo the classification of finite simple groups

to be closed under normal subgroups and homomorphic images (see Lemma 4(1)), but not closed under extensions in general (see [4, Ch. V, Example 2]). In [5, Theorem 3.5] and [6, Corollary 6] it is proven that, if $1 = G_0 < G_1 < \ldots < G_n = G$ is a composition series of G that is a refinement of a chief series, then G satisfies E_π if and only if $\mathrm{Aut}_G(G_i/G_{i-1})$ satisfies E_π for each $i = 1, \ldots, n$.

The class of C_π-groups is closed under extensions (see Lemma 5) but not closed under normal subgroups in general (see the example below). In the present paper, by using the classification of finite simple groups, we show that the class of C_π-groups is closed under homomorphic images (mod CFSG, see Lemma 9), and give a criterion for a finite group to satisfy C_π in terms of a normal series of the group. The main result is the following:

Theorem 1. (mod CFSG) *Let π be a set of primes, let H be a π-Hall subgroup, and A be a normal subgroup of a C_π-group G. Then $HA \in C_\pi$.*

Corollary 2. (a conjugacy criterion for Hall subgroups, mod CFSG). *Let π be a set of primes and let A be a normal subgroup of G. Then $G \in C_\pi$ if and only if $G/A \in C_\pi$ and, for a π-Hall*[2] *subgroup K/A of G/A its complete preimage K satisfies C_π. In particular, if $|G : A|$ is a π'-number, then $G \in C_\pi$ if and only if $A \in C_\pi$.*

Using the corollary, at the end of the paper we give an algorithm that reduces the problem whether a finite groups satisfies C_π to the same problem in some almost simple groups. In view of Theorem 1 notice that we do not know any counterexample to the following hypothesis.

Hypothesis 3. *Let π be a set of primes and let A be a (not necessary normal) subgroup of a finite C_π-group G containing a π-Hall subgroup of G. Then $A \in C_\pi$.*

In the hypothesis the condition that A includes a π-Hall subgroup of G cannot be weaken by the condition that the index of A is a π'-number. Indeed, consider $B_3(q) \simeq \mathrm{P}\Omega_7(q)$, where $q-1$ is divisible by 12 and is not divisible by 8 and 9. In view of [2, Lemma 6.2], $\mathrm{P}\Omega_7(q)$ is a $C_{\{2,3\}}$-group and its $\{2,3\}$-Hall subgroup is included in a monomial subgroup. On the other hand, $\Omega_7(2)$ is known to be isomorphic to a subgroup of $\mathrm{P}\Omega_7(q)$ and, under the above stated conditions on q, its index is not divisible by 2 and 3. However $\Omega_7(2)$ does not possess a $\{2,3\}$-Hall subgroup, i.e., it is not even an $E_{\{2,3\}}$-group.

1 Notations and preliminary results

By π we always denote a set of primes, and the term "group" always means a finite group.

The lemmas below are known and their proof do not refer to the classification of finite simple groups.

Lemma 4. [4, Ch. IV, (5.11), Ch. V, Theorem 3.7] *Let A be a normal subgroup of G. Then the following holds.*

(1) *If H is a π-Hall subgroup of G then $H \cap A$ is a π-Hall subgroup of A, while HA/A is a π-Hall subgroup of G/A.*

(2) *If all factors of a subnormal series of G are either π- or π'- groups then $G \in D_\pi$.*

---

[2] Since $G/A \in C_\pi$ the phrase "for a π-Hall subgroup K/A of G/A" in the statement can be interpreted as "for every" and "for some", and both of them are correct.

Notice that (2) of Lemma 4 follows from the famous Chunikhin's theorem on π-solvable groups and the Feit-Thompson Odd Order Theorem.

Lemma 5. (Chunikhin; see also [1, Theorems C1 and C2] or [4, Ch. V, (3.12)]) *Let A be a subnormal subgroup of G. If both A and G/A satisfy C_π, then $G \in C_\pi$.*

Lemma 6. [3, Lemma 2.1(e)] *Let A be a normal subgroup of G such that G/A is a π-group and let M be a π-Hall subgroup of A. Then a π-Hall subgroup H of G with $H \cap A = M$ exists if and only if G acting by conjugation leaves $\{M^a \mid a \in A\}$ invariant.*

EXAMPLE. Suppose that $\pi = \{2, 3\}$. Let $G = \mathrm{GL}_5(2) = \mathrm{SL}_5(2)$ be a group of order $99999360 = 2^{10} \cdot 3^2 \cdot 5 \cdot 7 \cdot 31$. Assume then that $\iota : x \in G \mapsto (x^t)^{-1}$ and $\widehat{G} = G \leftthreetimes \langle \iota \rangle$ is a natural semidirect product. By [7, Theorem 1.2], G possesses π-Hall subgroups and each π-Hall subgroup is the stabilizer of a series of subspaces $V = V_0 < V_1 < V_2 < V_3 = V$, where V is a natural module of G and $\dim V_k/V_{k-1} \in \{1, 2\}$ for every $k = 1, 2, 3$. Hence G possesses exactly three classes of conjugate π-Hall subgroups with representatives

$$H_1 = \begin{pmatrix} \boxed{\mathrm{GL}_2(2)} & & * \\ & \boxed{1} & \\ 0 & & \boxed{\mathrm{GL}_2(2)} \end{pmatrix},$$

$$H_2 = \begin{pmatrix} \boxed{1} & & * \\ & \boxed{\mathrm{GL}_2(2)} & \\ 0 & & \boxed{\mathrm{GL}_2(2)} \end{pmatrix}, \text{ and } H_3 = \begin{pmatrix} \boxed{\mathrm{GL}_2(2)} & & * \\ & \boxed{\mathrm{GL}_2(2)} & \\ 0 & & \boxed{1} \end{pmatrix}.$$

Notice that $N_G(H_k) = H_k$, $k = 1, 2, 3$, since H_k is parabolic. By Lemma 4(1), for every π-Hall subgroup H of $\widehat{G}$ the subgroup $H \cap G$ is conjugate to one of subgroups H_1, H_2, and H_3. The class containing H_1 is invariant under ι. Hence by Lemma 6 there exists a π-Hall subgroup H of $\widehat{G}$ with $H \cap G = H_1$. Moreover $H = N_{\widehat{G}}(H_1)$. The classes containing H_2 and H_3 are permuted by ι. So from Lemmas 4(1) and 6 it follows that these subgroups do not lie in π-Hall subgroups of $\widehat{G}$. Thus $\widehat{G}$ has exactly one class of conjugate π-Hall subgroups, and so satisfies C_π, in contrast to its normal subgroup G.

Lemma 7. *Let A be a normal subgroup and let H be a π-Hall subgroup of a C_π-group G. Then $N_G(HA)$ and $N_G(H \cap A)$ satisfy C_π.*

Proof. Note that both $N_G(HA)$ and $N_G(H \cap A)$ include H, and so they satisfy E_π. Let K be a π-Hall subgroup of $N_G(HA)$. Since $HA \trianglelefteq N_G(HA)$ and $|N_G(HA) : HA|$ is a π'-number, we have $K \leq HA$ and $KA = HA$. If $x \in G$ is chosen so that $K = H^x$ then $(HA)^x = H^x A = KA = HA$ and so $x \in N_G(HA)$. Therefore $N_G(HA) \in C_\pi$.

Suppose that K is a π-Hall subgroup of $N_G(H\cap A)$. Then $K(H\cap A)=K$ and $K\cap A=H\cap A$. If $x\in G$ is chosen so that $K=H^x$, then $(H\cap A)^x=H^x\cap A=K\cap A=H\cap A$ and so $x\in N_G(H\cap A)$. Therefore $N_G(H\cap A)\in C_\pi$. ☐

Lemma 8. ([6, Corollary 9] mod CFSG) *Each π-Hall of a homomorphic image of an E_π-group G is the image of a π-Hall subgroup of G.*

By Lemma 8 it is immediate that C_π is preserved under homomorphisms.

Lemma 9. (mod CFSG) *Let A be a normal subgroup of a C_π-group G. Then $G/A\in C_\pi$.*

Proof. Put $G/A=\overline{G}$. Since all π-Hall subgroups of G are conjugate, it is enough to show that for every π-Hall subgroup $\overline{K}$ of $\overline{G}$ there exists a π-Hall subgroup U of G such that $UA/A=\overline{K}$. The existence of U follows from Lemma 8. ☐

If G -is a group then a *G-class of π-Hall subgroups* is a class of conjugate π-Hall subgroups of G. Let A be a subnormal subgroup of a E_π-group G. A subgroup $H\cap A$ of A, where H is a π-Hall subgroup of G is called a *G-induced π-Hall subgroup* of A. Thus the set $\{(H\cap A)^a\mid a\in A\}$, where H is a π-Hall subgroup of G is called an *A-class of G-induced π-Hall subgroups.* We denote the number of all A-classes of G-induced π-Hall subgroups by $k_\pi^G(A)$. Let $k_\pi(G)=k_\pi^G(G)$ be the number of classes of π-Hall subgroups of G. Clearly $k_\pi^G(A)\leqslant k_\pi(A)$.

Recall that a finite group G is called *almost simple*, if G possesses a unique minimal normal subgroup S and S is a nonabelian finite group (equivalently, up to isomorphism, $S\simeq \mathrm{Inn}(S)\leq G\leq \mathrm{Aut}(S)$ for a nonabelian finite simple group S). The proof of Theorem 1 uses the following statement on the number of classes of π-Hall subgroups in finite simple groups.

Theorem 10. ([3, Theorem 1.1], mod CFSG) *Let π be a set of primes and G be an almost simple finite E_π-group with the (nonabelian simple) socle S. Then the following hold.*

(1) *If $2\notin\pi$ then $k_\pi^G(S)=1$.*

(2) *If $3\notin\pi$ then $k_\pi^G(S)\in\{1,2\}$.*

(3) *If $2,3\in\pi$ then $k_\pi^G(S)\in\{1,2,3,4,9\}$.*

In particular, $k_\pi^G(S)$ is a π-number..

Lemma 11. *Let H be a π-Hall subgroup, let A be a normal subgroup of G, and $HAC_G(A)$ is normal in G (this condition is satisfied, if $HA\trianglelefteq G$). Then an A-class of π-Hall subgroups is a class of G-induced π-Hall subgroups if and only if it is H-invariant.*

Proof. If K is a π-Hall subgroup of G then $K\leq HAC_G(A)$, and so $KAC_G(A)=HAC_G(A)$. Since the A-class $\{(K\cap A)^a\mid a\in A\}$ is K-invariant, it follows that it is invariant under $HAC_G(A)=KAC_G(A)$, and so under H.

Conversely, without loss of generality we may assume that $G=HA$ and the claim follows from Lemma 6. ☐

Lemma 12. *Let H be a π-Hall subgroup, let A be a normal subgroup of G, and $HA\trianglelefteq G$. Then $k_\pi^G(A)=k_\pi^{HA}(A)$.*

Proof. Since HA is a normal subgroup of G, each π-Hall subgroup of G lies in HA. Hence $k_\pi^G(A)=k_\pi^{HA}(A)$. ☐

Lemma 13. *Let H be a π-Hall subgroup, let A be a normal subgroup of G, and $HA \trianglelefteq G$. Then the following are equivalent:*
(1) $k_\pi^G(A) = 1$.
(2) $HA \in C_\pi$.
(3) *Every two π-Hall subgroups of G are conjugate by an element of A.*

Proof. (1) $\Rightarrow$ (2). If K is a π-Hall subgroup of HA then by (1) $H \cap A$ and $K \cap A$ are conjugate in A. We may assume that $H \cap A = K \cap A$. Then H and K lie in $N_{HA}(H \cap A)$. By the Frattini argument, $HA = N_{HA}(H \cap A)A$. So,

$$N_{HA}(H \cap A)/N_A(H \cap A) = N_{HA}(H \cap A)/N_{HA}(H \cap A) \cap A \simeq N_{HA}(H \cap A)A/A = HA/A$$

is a π-group. Thus $N_{HA}(H \cap A)$ possesses a normal series

$$N_{HA}(H \cap A) \geq N_A(H \cap A) \geq H \cap A \geq 1$$

such that every factor of the series is either a π- or π'- group, and, by Lemma 4(2), satisfies D_π. In particular, H and K are conjugate in $N_{HA}(H \cap A)$.

(2) $\Rightarrow$ (3) and (3) $\Rightarrow$ (1) are evident. □

Lemma 14. *Let H be a π-Hall subgroup, let $A = A_1 \times \ldots \times A_s$ be a normal subgroup of G, and $G = HAC_G(A)$. Then for every $i = 1, \ldots, s$ the following hold:*
(1) $N_G(A_i) = N_H(A_i)AC_G(A)$.
(2) *$N_H(A_i)$ is a π-Hall subgroup of $N_G(A_i)$.*
(3) $k_\pi^{\mathrm{Aut}_G(A_i)}(\mathrm{Inn}(A_i)) = k_\pi^{N_G(A_i)}(A_i)$.

Proof. (1) follows since $G = HAC_G(A)$ and $AC_G(A) \leq N_G(A_i)$. Using (1) and the identity $N_H(A_i) \cap AC_G(A) = H \cap AC_G(A)$, we see that $|N_G(A_i) : N_H(A_i)| = |AC_G(A) : (H \cap AC_G(A))|$ is a π'-number, whence (2).

Assume that $\rho : A_i \to \mathrm{Inn}(A_i)$ is the natural epimorphism. Since $\mathrm{Ker}(\rho) = Z(A_i)$ is an abelian group, the kernel of ρ possesses a unique π-Hall subgroup which lies in each π-Hall subgroup of A_i. Hence the map $H \mapsto H\rho$ defines a bijection between the sets of π-Hall subgroups of A_i and $\mathrm{Inn}(A_i)$, and also induces a bijection (we denote it by the same symbol σ) between the sets Δ and Γ of A_i- and $\mathrm{Inn}(A_i)$- classes of π-Hall subgroups, respectively. We show that the restriction of σ on the set Δ_0 of all A_i-classes of $N_G(A_i)$-induced π-Hall subgroups is a bijective map from Δ_0 onto the set Γ_0 of all $\mathrm{Inn}(A_i)$-classes of $\mathrm{Aut}_G(A_i)$-induced π-Hall subgroups. Since $A = A_1 \times \ldots \times A_s$; therefore, (1) implies $N_G(A_i) = N_H(A_i)A_iC_G(A_i)$. The normalizer $N_G(A_i)$ permutes elements from Δ acting by conjugation on the π-Hall subgroups of A_i. Thus, it acts on Δ. By Lemma 11, Δ_0 is the union of all one-element orbits under this action. By using σ, define an equivalent action of $N_G(A_i)$ on Γ. Since $C_G(A_i)$ lies in the kernel of both actions, the induced actions of $\mathrm{Aut}_G(A_i) = N_G(A_i)/C_G(A_i)$ on Δ and Γ are well-defined. It is easy to see that the action of $\mathrm{Aut}_G(A_i)$ on Γ defined in this way coincides with the natural action of the group on the set of $\mathrm{Inn}(A_i)$-classes of conjugate π-Hall subgroups. Since

$$\mathrm{Aut}_G(A_i)/\,\mathrm{Inn}(A_i) \simeq N_G(A_i)/A_iC_G(A_i) \simeq N_H(A_i)/(N_H(A_i) \cap A_iC_G(A_i))$$

is a π-group, by Lemma 11, Γ_0 coincides with the union of one-element orbits of $\mathrm{Aut}_G(A_i)$ on Γ. By the definition of the action, Γ_0 is the image of Δ_0 under σ. Since σ is a bijection,

$$k_\pi^{N_G(A_i)}(A_i) = |\Delta_0| = |\Gamma_0| = k_\pi^{\mathrm{Aut}_G(A_i)}(\mathrm{Inn}(A_i)).$$

(3) follows. □

Suppose $A = A_1 \times \cdots \times A_s$ and for every $i = 1, \ldots, s$ by $\mathcal{K}_i$ we denote an A_i-class of π-Hall subgroups of A_i. The set

$$\mathcal{K}_1 \times \cdots \times \mathcal{K}_s = \{\langle H_1, \ldots, H_s\rangle \mid H_i \in \mathcal{K}_i,\ i = 1, \ldots, s\}$$

is called the *product of classes* $\mathcal{K}_1, \ldots, \mathcal{K}_s$. Clearly $\mathcal{K}_1 \times \cdots \times \mathcal{K}_s$ is an A-class of π-Hall subgroups of A. It is also clear that for a normal subgroup A of G every A-class of G-induced π-Hall subgroups is a product of some A_1-, …, A_s- classes of G-induced π-Hall subgroups. In particular, $k_\pi^G(A_i) \leqslant k_\pi^G(A)$ for every $i = 1, \ldots, s$. The reverse inequality fails in general.

Lemma 15. *Let H be a π-Hal subgroupl and let $A = A_1 \times \cdots \times A_s$ be a normal subgroup of G. Assume also that the subgroups $A_1, \ldots, A_s$ are normal in G and $G = HAC_G(A)$. Then $k_\pi^G(A) = k_\pi^G(A_1) \cdot \ldots \cdot k_\pi^G(A_s)$.*

Proof. Two π-Hall subgroups P and Q of A are conjugate in A if and only if π-Hall subgroups $P \cap A_i$ and $Q \cap A_i$ of A_i are conjugate in A_i for every $i = 1, \ldots, s$. In order to prove the claim it is enough to show that the product of A_1-, …, A_s- classes of G-induced π-Hall subgroups is an A-class of G-induced π-Hall subgroups as well. Assume that $U_1, \ldots, U_s$ are G-induced π-Hall subgroups of $A_1, \ldots, A_s$, respectively. We show that

$$U = \langle U_1, \ldots, U_s\rangle = U_1 \times \ldots \times U_s$$

is a G-induced π-Hall subgroup of A. By Lemma 11 it is enough to show that for every $h \in H$ there exists $a \in A$ with $U^h = U^a$. Since $U_i = K_i \cap A_i$ for appropriate π-Hall subgroup K_i of G, the set $\left\{U_i^{x_i} \mid x_i \in A_i\right\}$ is K_i-invariant, so this set is invariant under $K_iA = HA$. In particular, $U_i^h = U_i^{a_i}$ for some $a_i \in A_i$. Thus

$$U^h = U_1^h \times \ldots \times U_s^h = U_1^{a_1} \times \ldots \times U_s^{a_s} = U_1^a \times \ldots \times U_s^a = U^a,$$

where $a = a_1 \ldots a_s \in A$. □

Lemma 16. *Let H be a π-Hall subgroup, let $A = A_1 \times \cdots \times A_s$ be a normal subgroup of G that acts transitively on the set $\{A_1, \ldots, A_s\}$ by conjugation, and $G = HAC_G(A)$. Then $k_\pi^G(A) = k_\pi^G(A_i) = k_\pi^{N_G(A_i)}(A_i)$ for every $i = 1, \ldots, s$.*

Proof. We show that each G-induced π-Hall subgroup of A_i is also a $N_G(A_i)$-induced subgroup. Indeed, if K is a π-Hall subgroup of G, then $G = KAC_G(A)$ and, by Lemma 14, the identity $N_G(A_i) = N_K(A_i)AC_G(A)$ holds. Moreover $K \cap A_i = N_K(A_i) \cap A_i$ and $N_K(A_i)$ is a π-Hall subgroup of $N_G(A_i)$. So every G-induced π-Hall subgroup of A_i is also a $N_G(A_i)$-induced π-Hall subgroup, in particular, $k_\pi^G(A_i) \leqslant k_\pi^{N_G(A_i)}(A_i)$.

Now we show that if $x, y \in G$ are in the same coset of G by $N_G(A_1)$, then for every G-induced π-Hall subgroup U_1 of A_1 the subgroups U_1^x and U_1^y are conjugate in $A_i = A_1^x = A_1^y$. It is enough to show that the subgroups U_1 and U_1^t, where $t = xy^{-1} \in N_G(A_1)$, are conjugate in A_1. Put $t = hac$, $a \in A$, $c \in C_G(A)$, $h \in H$. Since U_1^h and U_1^{hac} are conjugate in A_1, it is enough to show that U_1 and U_1^h are conjugate in A_1. Since $(ac)^{h^{-1}} \in AC_G(A) \leq N_G(A_1)$, the element h normalizes A_1 as well. Assume that $U_1 = U \cap A_1$ for a G-induced π-Hall subgroup U of A. Suppose that $\mathcal{K}$ is an A-class of π-Hall subgroups containing U, and let $\mathcal{K} = \mathcal{K}_1 \times \cdots \times \mathcal{K}_s$, where $\mathcal{K}_i$ is an A_i-class of G-induced π-Hall subgroups. Clearly, $U_1 \in \mathcal{K}_1$. Since, by Lemma 11, $\mathcal{K}$ is H-invariant, H acts on the set

$\{\mathcal{K}_1, \dots, \mathcal{K}_s\}$. The element h normalizes A_1, hence it fix the A_1-class $\mathcal{K}_1$. In particular, the subgroups U_1 and U_1^h are in $\mathcal{K}_1$, and so they are conjugate in A_1.

Assume that $f \in H$ and $A_1^f = A_i$. For an A_1-class of π-Hall subgroups $\mathcal{K}_1$ we define the A_i-class $\mathcal{K}_1^f$ by $\mathcal{K}_1^f = \{U_1^f \mid U_1 \in \mathcal{K}_1\}$. As we noted above, $\mathcal{K}_1^f$ is an A_i-class of G-induced π-Hall subgroups.

Let $h_1 = 1, h_2 \dots, h_s$ be the right transversal of $N_H(A_1)$ in H. Since G acts transitively, up to renumbering we may assume that $A_i = A_1^{h_i}$ and $(N_G(A_1))^{h_i} = N_G(A_i)$. So $k_\pi^{N_G(A_1)}(A_1) = k_\pi^{N_G(A_i)}(A_i)$ for $i = 1, \dots, s$. Consider

$$\sigma : \mathcal{K}_1 \mapsto \mathcal{K}_1^{h_1} \times \cdots \times \mathcal{K}_1^{h_s},$$

mapping an A_1-class of $N_G(A_1)$-induced π-Hall subgroups $\mathcal{K}_1$ to an A-class of π-Hall subgroups. Note that $\mathcal{K}_1^{h_1} \times \dots \times \mathcal{K}_1^{h_s}$ is always H-invariant and, by Lemma 11, it is an A-class of G-induced π-Hall subgroups. Notice also that σ is injective, and so $k_\pi^{N_G(A_1)}(A_1) \leqslant k_\pi^G(A)$. Consider the restriction τ of σ on the set of A_1-classes of G-induced π-Hall subgroups. We need to show that the image of τ coincides with the set of A-classes of G-induced π-Hall subgroups in order to complete the prove, since in such case we derive the inequality $k_\pi^G(A_1) \geqslant k_\pi^G(A)$.

Let $\mathcal{K} = \mathcal{K}_1 \times \cdots \times \mathcal{K}_s$ be an A-class of G-induced π-Hall subgroups. It is enough to show that $\mathcal{K}_i = \mathcal{K}_1^{h_i}$ for every $i = 1, \dots, s$. Since G acts transitively on the set $\{A_1, \dots, A_s\}$, there exists an element $g \in G$ such that $A_1^g = A_i$. Let gct, where $a \in A$, $c \in C_G(A)$, and $t \in H$. Then $A_1^t = A_i$ and $t \in N_H(A_1)h_i$. As we already proved, $\mathcal{K}_1^t = \mathcal{K}_1^{h_i}$. By Lemma 11, $\mathcal{K}$ is H-invariant. Hence $\mathcal{K}_i = \mathcal{K}_1^t = \mathcal{K}_1^{h_i}$ and $\mathcal{K} = \mathcal{K}_1^{h_1} \times \cdots \times \mathcal{K}_1^{h_s} = \mathcal{K}_1\tau$. $\square$

2 A conjugacy criterion for Hall subgroups

In the section we prove Theorem 1, Corollary 2, and provide an algorithm for determining whether G satisfies C_π by using a normal series of G.

Proof of Theorem 1. Assume that the claim is not true, and let G be a counterexample of minimal order. Then G possesses a π-Hall subgroup H and a normal subgroup A such that HA does not satisfy C_π. We choose A to be minimal. Let K be a π-Hall subgroup of HA that is not conjugate with H in HA. We divide into several steps the process of canceling the group G.

Clearly

(1) $HA = KA$.

(2) *A is a minimal normal subgroup of G.*

Otherwise assume that M is a nontrivial normal subgroup of G that is contained properly in A. Put $\overline{G} = G/M$ and, given a subgroup B of G, denote BM/M by $\overline{B}$. By Lemma 9, $\overline{G}$ satisfies C_π, $\overline{H}$ and $\overline{K}$ are π-Hall subgroups of $\overline{G}$, $\overline{A}$ is a normal subgroup of $\overline{G}$, $\overline{HA} = \overline{KA}$ and $|\overline{G}| < |G|$. In view of the minimality of G, the group $\overline{HA}$ satisfies C_π. So $\overline{H}$ and $\overline{K}$ are conjugate by an element of $\overline{A}$. Hence, the subgroups HM and KM are conjugate by an element of A. Without loss of generality, we may assume that $HM = KM$. In view of the choice of A, the group HM satisfies C_π. Hence H and K are conjugate by an element of $M \leq A$; a contradiction.

(3) $A \notin C_\pi$. *In particular, A is not solvable.*

Otherwise, by Lemma 5, the group HA satisfies C_π as an extension of a C_π-groups by a π-group.

(4) *HA is a normal subgroup of G.*

Otherwise $N_G(HA)$ is a proper subgroup of G and, by Lemma 7, we have $N_G(HA) \in C_\pi$. In view of the minimality of G, it follows that $HA \in C_\pi$; a contradiction.

In view of (2) and (3)

(5) *A is a direct product of simple nonabelian groups $S_1, \dots, S_m$. The group G acts transitively on the set $\Omega = \{S_1, \dots, S_m\}$ by conjugation.*

Let $\Delta_1, \dots, \Delta_s$ be the orbits of HA on Ω, and put $T_j = \langle \Delta_j \rangle$ for every $j = 1, \dots, s$. In view of (4) and (5),

(6) *G acts transitively on $\{T_1, \dots, T_s\}$ by conjugation. The subgroup A is a direct product of $T_1, \dots, T_s$ and each of these subgroups is normal in HA.*

Assume that $S \in \Omega$, and let T be a subgroup, generated by the orbit from the set $\{\Delta_1, \dots, \Delta_s\}$ that contains S. By Lemma 16,

(7) $k_\pi^{HA}(T) = k_\pi^{HA}(S)$.

By (7) and Lemmas 12 and 15

(8) $k_\pi^G(A) = k_\pi^{HA}(A)\big(k_\pi^{HA}(T)\big)^s = \big(k_\pi^{HA}(S)\big)^s$.

By (8), Theorem 10 and Lemmas 14 and 16

(9) *$k_\pi^G(A)$ is a π-number.*

By Lemma 11,

(10) *HA fixes every A-class of G-induced π-Hall subgroups.*

Since $G \in C_\pi$,

(11) *G acts transitively on the set of A-classes of G-induced π-Hall subgroups.*

In view of (10), the subgroup HA lies in the kernel of this action. Now, by (11),

(12) *$k_\pi^G(A)$ is a π'-number.*

By (9) and (12), it follows that

(13) $k_\pi^G(A) = 1$.

Now by Lemma 13,

(14) *$HA \in C_\pi$; a contradiction.* ☐

Proof of Corollary 2. Necessity. If $G \in C_\pi$, then, by Lemma 9, $G/A \in C_\pi$ as well. Let K/A be a π-Hall subgroup of G/A. By Lemma 8, there exists a π-Hall subgroup H of G such that $K = HA$. By Theorem 1, the group $K = HA$ satisfies C_π.

Sufficiency. Let H be a π-Hall subgroup of K. Since K/A is a π-Hall subgroup of G/A, we have $|G : H| = |G : K| \cdot |K : H|$ is a π'-number, and so H is a π-Hall subgroup of G. In particular, $G \in E_\pi$. Let H_1, H_2 be π-Hall subgroups of G. Since $G/A \in C_\pi$, the subgroups H_1A/A and H_2A/A are conjugate in G/A, and we may assume that $H_1A = H_2A$. However $H_1A \in C_\pi$, and so H_1 and H_2 are conjugate. Thus $G \in C_\pi$. ☐

Lemma 17. *Assume that $G = HA$, where H is a π-Hall subgroup, A is a normal subgroup of G, and $A = S_1 \times \ldots \times S_k$ is a direct product of simple groups. Then $G \in C_\pi$ if and only if $\mathrm{Aut}_G(S_i) \in C_\pi$ for every $i = 1, \dots, k$.*

Proof. By the Hall theorem and Lemma 5 we may assume that $S_1, \dots, S_k$ are nonabelian simple groups, and so the set $\{S_1, \dots, S_k\}$ is invariant under the action of G by conjugation. Moreover this set is partitioned into the orbits $\Omega_1, \dots, \Omega_m$. Denote an element of Ω_j by S_{i_j}. By Lemmas 15 and 16, we obtain

$$k_\pi^G(A) = k_\pi^G(S_{i_1}) \cdot \ldots \cdot k_\pi^G(S_{i_m}). \tag{1}$$

Lemmas 12 and 13 imply that $HA \in C_\pi$ if and only if $k_\pi^G(A) = 1$, and in view of (1), if and only if $k_\pi^G(S_i) = 1$ for every i. By Lemmas 14 and 16, $k_\pi^G(S_i) = k_\pi^{N_G(S_i)}(S_i) = k_\pi^{\mathrm{Aut}_G(S_i)}(S_i)$ for every i. Moreover, since $|\operatorname{Aut}_G(S_i) : S_i|$ is a π-number, in view of Lemma 13, $k_\pi^{\mathrm{Aut}_G(S_i)}(S_i) = 1$ holds if and only if $\operatorname{Aut}_G(S_i) \in C_\pi$. Therefore, $k_\pi^G(S_i) = 1$ for every i if and only if $\operatorname{Aut}_G(S_i) \in C_\pi$. □

Now we are able to give an algorithm reducing the problem, whether a finite group satisfies C_π, to the check of C_π-property in some almost simple groups. Assume that

$$G = G_0 > G_1 > \ldots > G_n = 1 \tag{2}$$

is a chief series of G. Put $H_1 = G = G_0$. Suppose that for some $i = 1, \ldots, n$ the group H_i is constructed so that $G_{i-1} \leq H_i$ and H_i/G_{i-1} is a π-Hall subgroup of G/G_{i-1}. Since (2) is a chief series,

$$G_{i-1}/G_i = S_1^i \times \ldots \times S_{k_i}^i,$$

where $S_1^i, \ldots, S_{k_i}^i$ are simple groups. We check whether

$$\operatorname{Aut}_{H_i}(S_1^i) \in C_\pi, \ldots, \operatorname{Aut}_{H_i}(S_{k_i}^i) \in C_\pi.$$

If so then by Lemma 17, we have $H_i/G_i \in C_\pi$ and we can take a complete preimage of a π-Hall subgroup of H_i/G_i to be equal to H_{i+1}. Otherwise, by Corollary 2, we deduce that $G \notin C_\pi$ and stop the process. By Corollary 2 it follows, that G satisfies C_π if and only if the group H_{n+1} can be constructed. Notice that in this case H_{n+1} is a π-Hall subgroup of G.

Corollary 18. *If either $2 \notin \pi$ or $3 \notin \pi$, then $G \in C_\pi$ if and only if every nonabelian composition factor of G satisfies C_π.*

Proof. The *sufficiency* follows from Lemma 5. We prove the *necessity.* By the above algorithm, we may assume that $S \leq G \leq \operatorname{Aut}(S)$ for a nonabelian finite simple group S, and G/S is a π-number. We need to show that $S \in C_\pi$. Assume the contrary. Then Lemmas 11 and 13 imply that G stabilizes precisely one class of π-Hall subgroups of S. Therefore, S possesses at least three classes of π-Hall subgroups. On the other hand, since either $2 \notin \pi$, or $3 \notin \pi$, by Theorem 10 the number of classes of π-Hall subgroups in S is not greater than 2; a contradiction. □

Notice that in the case $2 \notin \pi$ the corollary is immediate from [8, Theorem A] and Lemma 5.

References

[1] P.Hall, Theorems like Sylow‘s, Proc.London Math.Soc., 6, N 22 (1956), 286–304.

[2] D.O.Revin, E.P.Vdovin, Hall subgroups of finite groups, *Ischia Group Theory 2004: Proceedings of a Conference in Honor of Marcel Herzog*, Contemporary Mathematics, AMS, 402 (2006), 229–263.

[3] D.O.Revin, E.P.Vdovin, On the number of classes of conjugate Hall subgroups in finite simple groups, *J.Algebra*, (to appear). (Available at http://arxiv.org/abs/0912.1922).

[4] M.Suzuki, *Group Theory* II. New York, Springer-Verl. 1986.

[5] F.Gross, On the existence of Hall Subgroups, *J.Algebra*, **98** (1986), N1, 1–13.

[6] D.O.Revin, E.P.Vdovin. Existence criterion for Hall subgroups of finite groups, *J.Gr.Theory* (to appear). (Available at http://arxiv.org/abs/0803.3868).

[7] D.O.Revin, Hall π-subgroups of finite Chevalley groups whose characteristic belongs to π, *Sib.Adv.Math.*, 9, N 2 (1999), 25–71.

[8] F.Gross, Conjugacy of odd order Hall subgroups, *Bull.London Math.Soc.*, **19** (4) (1987), N 79, 311-319.